

Ovarian Cancer™

U P D A T E

Conversations with Oncology Investigators
Bridging the Gap between Research and Patient Care

EDITOR

Neil Love, MD

INTERVIEWS

Tate Thigpen, MD

Ursula A Matulonis, MD

Robert A Burger, MD

Bradley J Monk, MD



Ovarian Cancer Update

A Continuing Medical Education Audio Series

OVERVIEW OF ACTIVITY

Optimal oncologic management of ovarian cancer begins with intensive surgical staging and cytoreduction, followed by primary chemotherapy and, for most patients, subsequent medical management when platinum-resistant relapsed disease prevails. Although many single-agent and combination cytotoxic recurrence regimens have been studied, only recently has the advent of antibody and small-molecule growth-inhibitory targeted agents been integrated into the ovarian cancer research milieu. It is hoped that the results from these trials will lead to the emergence of new therapeutic agents and changes or enhancements in the indications for existing treatment strategies, ultimately improving the duration and quality of life for patients with metastatic ovarian cancer. In order to offer optimal care to the ovarian cancer population — including the option of clinical trial participation — practicing oncologists must be well informed of these advances. By providing access to the latest research developments and expert perspectives through one-on-one discussion with leading investigators, *Ovarian Cancer Update* will assist medical and gynecologic oncologists with the formulation of up-to-date clinical management strategies.

LEARNING OBJECTIVES

- Recognize the contributory roles of the specialized interdisciplinary team in achieving best-practice surgical and medical outcomes for patients with ovarian cancer.
- Compare and contrast the risks and benefits of intraperitoneal and intravenous chemotherapy regimens when devising management strategies for patients with optimally debulked Stage II and Stage III ovarian cancer.
- Develop an evidence-based algorithm for the systemic treatment of recurrent platinum-sensitive and platinum-resistant ovarian cancer that optimizes long-term patient outcome and quality of life.
- Summarize the existing data and ongoing clinical trials focused on angiogenesis inhibition of ovarian cancer, and identify patients who may benefit from this therapeutic approach.
- Consider the utility of evaluating CA125 serum levels and radiographic monitoring of patients with ovarian cancer that is in a state of remission.
- Understand the potential role of the risk-adapted use of maintenance systemic therapy for patients with ovarian cancer who have demonstrated an initial treatment response.
- Recall the scientific rationale for molecular-targeted agents under active investigation for the treatment of ovarian cancer.
- Counsel appropriately selected patients with ovarian cancer about the availability of and participation in ongoing clinical trials.

ACCREDITATION STATEMENT

Research To Practice is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

CREDIT DESIGNATION STATEMENT

Research To Practice designates this educational activity for a maximum of 3 *AMA PRA Category 1 Credits™*. Physicians should only claim credit commensurate with the extent of their participation in the activity.

HOW TO USE THIS CME ACTIVITY

This CME activity contains both audio and print components. To receive credit, the participant should review the CME information, listen to the CDs and complete the Post-test and Educational Assessment and Credit Form located in the back of this monograph or on our website at ResearchToPractice.com/OCU. This monograph contains edited comments, clinical trial schemas, graphics and references that supplement the audio program. ResearchToPractice.com/OCU includes an easy-to-use, interactive version of this monograph with links to relevant full-text articles, abstracts, trial information and other web resources indicated here in [blue underlined text](#).

This program is supported by educational grants from Eli Lilly and Company, Genentech BioOncology and Ortho Biotech Products LP.

3 INTERVIEWS

Tate Thigpen, MD

Professor of Medicine
Director of Medical Oncology
University of MS Medical Center
Jackson, Mississippi

7 Ursula A Matulonis, MD

Medical Director and Program Leader
Gynecologic Oncology Program
Assistant Professor of Medicine
Harvard Medical School
Boston, Massachusetts

11 Robert A Burger, MD

Associate Professor of Clinical Obstetrics and Gynecology, Step II
Department of Obstetrics and Gynecology
Division of Gynecologic Oncology
University of California, Irvine Medical Center
Orange, California

14 Bradley J Monk, MD

Associate Professor and Director of Research
Division of Gynecologic Oncology
Chao Family Comprehensive Cancer Center
University of California, Irvine Medical Center
Orange, California

18 POST-TEST

19 EDUCATIONAL ASSESSMENT AND CREDIT FORM

If you would like to discontinue your complimentary subscription to *Ovarian Cancer Update*, please email us at Info@ResearchToPractice.com, call us at (800) 648-8654 or fax us at (305) 377-9998. Please include your full name and address, and we will remove you from the mailing list.

CONTENT VALIDATION AND DISCLOSURES

Research To Practice (RTP) is committed to providing its participants with high-quality, unbiased and state-of-the-art education. We assess potential conflicts of interest with faculty, planners and managers of CME activities. Real or apparent conflicts of interest are identified and resolved through a conflict of interest resolution process. In addition, all activity content is reviewed by both a member of the RTP scientific staff and an external, independent physician reviewer for fair balance, scientific objectivity of studies referenced and patient care recommendations.

FACULTY — The following faculty (and their spouses/partners) reported real or apparent conflicts of interest, which have been resolved through a conflict of interest resolution process: **Dr Thigpen** — Advisory Committee: Bristol-Myers Squibb Company, Eisai Inc, Genentech BioOncology, GlaxoSmithKline, Millennium Pharmaceuticals Inc, Novartis Pharmaceuticals Corporation, Ortho Biotech Products LP, Pfizer Inc, Sanofi-Aventis; Consulting Agreements: Abraxis BioScience, Amgen Inc, AstraZeneca Pharmaceuticals LP, Eli Lilly and Company; Speakers Bureau: Eli Lilly and Company, GlaxoSmithKline, Ortho Biotech Products LP. **Dr Matulonis** — Advisory Committee: Genentech BioOncology; Speakers Bureau: Amgen Inc. **Dr Burger** — Consulting Agreement: Eli Lilly and Company. **Dr Monk** — Paid Research: Eli Lilly and Company, GlaxoSmithKline, Ortho Biotech Products LP, Sanofi-Aventis; Speakers Bureau: Eli Lilly and Company, GlaxoSmithKline, Ortho Biotech Products LP.

EDITOR — **Dr Love** does not receive any direct remuneration from industry. Research To Practice receives funds in the form of educational grants to develop CME activities from the following commercial interests: Abraxis BioScience, AstraZeneca Pharmaceuticals LP, Aureon Laboratories Inc, Bayer Pharmaceuticals Corporation/Onyx Pharmaceuticals Inc, Biogen Idec, Bristol-Myers Squibb Company, Celgene Corporation, Eisai Inc, Eli Lilly and Company, Genentech BioOncology, Genomic Health Inc, GlaxoSmithKline, ImClone Systems Incorporated, Merck and Company Inc, Millennium Pharmaceuticals Inc, Novartis Pharmaceuticals Corporation, Ortho Biotech Products LP, OSI Oncology, Pfizer Inc, Roche Laboratories Inc, Sanofi-Aventis, Synta Pharmaceuticals Corp and Wyeth.

RESEARCH TO PRACTICE STAFF AND EXTERNAL REVIEWERS — The scientific staff and reviewers for Research To Practice have no real or apparent conflicts of interest to disclose.

This educational activity contains discussion of published and/or investigational uses of agents that are not indicated by the Food and Drug Administration. Research To Practice does not recommend the use of any agent outside of the labeled indications. Please refer to the official prescribing information for each product for discussion of approved indications, contraindications and warnings. The opinions expressed are those of the presenters and are not to be construed as those of the publisher or grantors.

Ovarian Cancer Update Downloadable Audio and Podcasts



Ovarian Cancer Update is available in MP3 format or as a Podcast delivered directly to your computer. To download complimentary copies of *OCU* or to subscribe to our free Podcasts, please visit www.ResearchToPractice.com/OCU.

What is a Podcast? Podcasts are audio files that are automatically delivered to Podcasting software on your computer, such as iTunes® or Juice Receiver, each time a new issue is available. You can listen to these files on your computer, or they can be quickly and easily transferred to your iPod® or other portable audio MP3 player for listening on the road, at home or while you exercise.

Please note that all of our other audio series are also available in these formats, and you may subscribe to as many Podcasts as you wish.



INTERVIEW

Tate Thigpen, MD

Dr Thigpen is Professor of Medicine and Director of Medical Oncology at the University of MS Medical Center in Jackson, Mississippi.

Tracks 1-12

- | | | | |
|----------------|---|-----------------|---|
| Track 1 | Conventional paclitaxel/carboplatin versus dose-dense weekly paclitaxel and carboplatin in advanced epithelial ovarian cancer, fallopian tube cancer (FTC) or primary peritoneal cancer (PPC) | Track 7 | Studies evaluating bevacizumab in combination with chemotherapy for patients with ovarian cancer |
| Track 2 | GOG-0126Q: A Phase II evaluation of pemetrexed in recurrent or persistent platinum-resistant ovarian cancer or PPC | Track 8 | Randomized studies of doublet chemotherapy versus single-agent platinum in platinum-sensitive, recurrent ovarian cancer |
| Track 3 | Perspective on the NCI clinical alert regarding intraperitoneal therapy for small-volume residual advanced ovarian cancer | Track 9 | Treatment algorithm for patients with platinum-resistant advanced ovarian cancer |
| Track 4 | Proposed GOG trial comparing intravenous to intraperitoneal therapy | Track 10 | Tolerability and side effects of agents commonly used for platinum-resistant advanced ovarian cancer |
| Track 5 | Ongoing GOG studies evaluating maintenance therapy in advanced ovarian cancer | Track 11 | Clinical investigations of novel agents in ovarian cancer — pertuzumab, enzastaurin and patupilone |
| Track 6 | Efficacy of bevacizumab during earlier-line therapy for advanced ovarian cancer | Track 12 | Increasing therapeutic options for recurrent ovarian cancer |

Select Excerpts from the Interview

Track 1

► **DR LOVE:** What important data sets in ovarian cancer were presented at ASCO this year?

► **DR THIGPEN:** The most interesting abstract was the Japanese study evaluating weekly paclitaxel with every three-week carboplatin versus every three-week carboplatin/paclitaxel in women with advanced ovarian cancer (Isonishi 2008). The response rate was essentially equivalent — approximately 55 percent — for both arms, but the median progression-free survival was 28

months versus 17 months in the weekly and every three-week arms, respectively, which is a dramatic difference. There were not enough events for a final survival analysis, but two-year survival was 83 percent with weekly paclitaxel and 77 percent with the every three-week regimen, which was statistically significant.

These data fly in the face of data reported six years ago from a study that evaluated weekly versus every three-week single-agent paclitaxel as second-line therapy, which revealed no difference between the two schedules (Rosenberg 2002). However, the Japanese data do correspond with the observations in breast cancer that weekly paclitaxel may be the most effective schedule (Sparano 2008).

The Japanese study needs to be repeated before we can routinely introduce the regimen into clinical practice, and the Gynecologic Oncology Group is planning on conducting that study in the near future.

Tracks 3-4

▶ **DR LOVE:** What are your thoughts about intraperitoneal (IP) therapy for ovarian cancer?

▶ **DR THIGPEN:** This is a controversial topic. In January 2006, the National Cancer Institute declared a combination of IP therapy and intravenous (IV) therapy to be the standard treatment for small-volume residual advanced ovarian cancer. However, they couldn't recommend a specific regimen (NCI 2006).

At our institution, IP therapy remains an experimental technique that is generally not used outside of a clinical trial for several reasons. First, it's quite toxic and many patients are unable to tolerate it. In the last GOG study of IP therapy, 48 percent of the patients received three or fewer cycles of IP therapy and only 42 percent were able to complete six cycles (Armstrong 2006).

Second, we are not certain whether the addition of IP therapy is superior to IV carboplatin/paclitaxel. In all of the Phase III studies of IP therapy conducted in the United States to date, the control arm has been an IV cisplatin-based regimen. Yet evidence from the last GOG study comparing IV cisplatin/paclitaxel to IV carboplatin/paclitaxel showed that the latter might be superior (Ozols 2003).

To address this question, GOG is planning to begin a trial in the next six to 12 months comparing a control arm of IV carboplatin/paclitaxel to two IP regimens. One IP regimen will utilize cisplatin and the other carboplatin.

Track 7

▶ **DR LOVE:** What studies have been conducted with bevacizumab for the treatment of advanced ovarian cancer?

▶ **DR THIGPEN:** The initial Phase II study, GOG-0170D, evaluated bevac-

zumab at 15 mg/kg every three weeks in patients who had received one or two prior chemotherapy regimens. The trial reported an objective response rate of 21 percent, a complete response rate of three percent and progression-free survival of at least six months in 40 percent (Burger 2007; [3.1, page 12]).

Two other Phase II studies evaluated bevacizumab in patients with more heavily pretreated disease. Among those patients who had received as many as three to five prior regimens, an increase in the complication rate was observed, specifically bowel complications (Cannistra 2007; Monk 2006). This suggests that if we use bevacizumab in ovarian cancer, it ought to be in the front-line setting or at first relapse.

- ▶ **DR LOVE:** What about bevacizumab combined with chemotherapy?
- ▶ **DR THIGPEN:** A trial evaluating cyclophosphamide with bevacizumab showed activity, but it was not a randomized study so we don't know whether the combination was better (Garcia 2008; [1.1]). Most of us feel strongly that administering bevacizumab with chemotherapy will be a better approach because of the possibility that it will improve the delivery of chemotherapy to the tumor by stabilizing the vasculature.

1.1

Phase II Trial of Bevacizumab and Low-Dose Metronomic Oral Cyclophosphamide in Recurrent Ovarian Cancer

“Our data suggest that the combination of bevacizumab and MC [metronomic chemotherapy] has significant activity in recurrent OC [ovarian cancer]. This was a population that was resistant to at least one line of platinum therapy having all progressed fewer than 12 months from a prior platinum therapy. The encouraging activity, time to progression, and median survival compare favorably with both conventional and investigational agents.”

SOURCE: Garcia AA et al. *J Clin Oncol* 2008;26(1):76-82. [Abstract](#)

Tracks 8-10

- ▶ **DR LOVE:** For patients with recurrent ovarian cancer, how do you define and approach platinum-sensitive versus platinum-resistant disease?
- ▶ **DR THIGPEN:** Chemosensitive disease is defined as that which responded to the previous treatment and did not progress for at least six months. Patients with chemosensitive disease should be treated with a regimen that is the same or similar to the initial one because their chances of responding are good. Disease that does not meet those criteria should be labeled as chemoresistant and should be treated with alternative drugs. For practical purposes, we talk about *platinum*-resistant or *platinum*-sensitive disease because all patients initially receive a platinum-based regimen. So patients who initially received paclitaxel/carboplatin and experience relapse more than six months later will fare better with a platinum-based regimen more so than anything else.

► **DR LOVE:** How do you treat platinum-resistant, recurrent ovarian cancer?

► **DR THIGPEN:** I prefer single agents in this setting because we don't have any proof that combinations are better. The six drugs that are commercially available, reimbursable and clearly have activity in platinum-resistant disease are pegylated liposomal doxorubicin (PLD), topotecan, gemcitabine, oral etoposide, weekly paclitaxel (if the patient received every three-week paclitaxel up front) and docetaxel.

All of these agents have response rates in the range of 15 to 20 percent and offer some survival benefit in the second-line setting. I generally start with PLD because it's administered once every four weeks, which is convenient for our patients. I use topotecan second, gemcitabine third, etoposide fourth, weekly paclitaxel fifth, and docetaxel sixth. Most of our patients will receive five or six different regimens.

There are two other agents that we should consider adding to that list — pemetrexed and bevacizumab. Bevacizumab, however, still has a question mark because of the increased complication rate among patients who have received multiple prior treatments. ■

SELECT PUBLICATIONS

Armstrong DK et al; Gynecologic Oncology Group. **Intraperitoneal cisplatin and paclitaxel in ovarian cancer.** *N Engl J Med* 2006;354(1):34-43. [Abstract](#)

Burger RA et al. **Phase II trial of bevacizumab in persistent or recurrent epithelial ovarian cancer or primary peritoneal cancer: A Gynecologic Oncology Group study.** *J Clin Oncol* 2007;25(33):5165-71. [Abstract](#)

Cannistra SA et al. **Phase II study of bevacizumab in patients with platinum-resistant ovarian cancer or peritoneal serous cancer.** *J Clin Oncol* 2007;25(33):5180-6. [Abstract](#)

Garcia AA et al. **Phase II clinical trial of bevacizumab and low-dose metronomic oral cyclophosphamide in recurrent ovarian cancer: A trial of the California, Chicago, and Princess Margaret Hospital phase II consortia.** *J Clin Oncol* 2008;26(1):76-82. [Abstract](#)

Isonishi S et al. **Randomized phase III trial of conventional paclitaxel and carboplatin (c-TC) versus dose dense weekly paclitaxel and carboplatin (dd-TC) in women with advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer: Japanese Gynecologic Oncology.** *Proc ASCO* 2008;[Abstract 5506](#).

Monk BJ et al. **Salvage bevacizumab (rhuMAB VEGF)-based therapy after multiple prior cytotoxic regimens in advanced refractory epithelial ovarian cancer.** *Gynecol Oncol* 2006;102(2):140-4. [Abstract](#)

National Cancer Institute (NCI). **NCI clinical announcement on intraperitoneal chemotherapy in ovarian cancer (January 5, 2006).** Available at: http://ctep.cancer.gov/highlights/clin_annc_010506.pdf. Accessed November 10, 2008.

Ozols RF et al; Gynecologic Oncology Group. **Phase III trial of carboplatin and paclitaxel compared with cisplatin and paclitaxel in patients with optimally resected stage III ovarian cancer: A Gynecologic Oncology Group study.** *J Clin Oncol* 2003;21(17):3194-200. [Abstract](#)

Rosenberg P et al. **Randomized trial of single agent paclitaxel given weekly versus every three weeks and with peroral versus intravenous steroid premedication to patients with ovarian cancer previously treated with platinum.** *Acta Oncol* 2002;41(5):418-24. [Abstract](#)

Sparano JA et al. **Weekly paclitaxel in the adjuvant treatment of breast cancer.** *N Engl J Med* 2008;358(16):1663-71. [Abstract](#)



INTERVIEW

Ursula A Matulonis, MD

Dr Matulonis is Medical Director and Program Leader of the Gynecologic Oncology Program and Assistant Professor of Medicine at Harvard Medical School in Boston, Massachusetts.

Tracks 1-18

- Track 1** GOG-0218: Carboplatin/paclitaxel with or without concurrent bevacizumab, with or without extended bevacizumab as up-front therapy for Stage III/IV epithelial ovarian cancer, PPC or FTC
- Track 2** Rationale for maintenance bevacizumab/erlotinib in optimally debulked ovarian cancer
- Track 3** Bevacizumab-associated bowel perforation in newly diagnosed versus recurrent ovarian cancer
- Track 4** Clinical activity of bevacizumab in chemotherapy-refractory, recurrent ovarian cancer
- Track 5** Bevacizumab and the palliation of symptomatic ascites in refractory ovarian cancer
- Track 6** Clinical use of bevacizumab for recurrent ovarian cancer
- Track 7** Bevacizumab-associated hypertension
- Track 8** Pan-VEGFR tyrosine kinase inhibitor cediranib in recurrent epithelial ovarian cancer
- Track 9** Identifying predictors of response to pertuzumab, a monoclonal antibody targeting the extracellular dimerization domain of the HER2 tyrosine kinase receptor
- Track 10** Intraperitoneal administration of chemotherapy in optimally debulked Stage III ovarian cancer
- Track 11** Therapeutic algorithm for patients with recurrent ovarian cancer
- Track 12** Gemcitabine/carboplatin in platinum-sensitive, recurrent ovarian cancer
- Track 13** Hypersensitivity reactions to carboplatin or paclitaxel
- Track 14** Treatment options for patients with platinum-resistant ovarian cancer
- Track 15** Potential role for poly(ADP-ribose) polymerase (PARP) inhibitors in ovarian cancer
- Track 16** Circulating tumor cells in ovarian cancer
- Track 17** Treatment approach for asymptomatic patients with rising CA125 concentrations on follow-up for ovarian cancer
- Track 18** Clinical applications of hormonal therapy in ovarian cancer

Select Excerpts from the Interview

Track 2

► **DR LOVE:** What is your treatment approach for optimally debulked ovarian cancer?

► **DR MATULONIS:** At the Dana-Farber/Harvard Cancer Center, patients with one centimeter or less of residual tumor after surgery either receive standard therapy — carboplatin/paclitaxel or IP cisplatin/paclitaxel — or enroll on a clinical trial.

We are conducting a pilot study evaluating IP carboplatin/paclitaxel with IV bevacizumab in women with newly diagnosed, optimally cytoreduced carcinoma of Müllerian origin. We also have a Phase II trial in which patients receive IV carboplatin/paclitaxel and bevacizumab for six months and then are randomly assigned to receive bevacizumab alone or bevacizumab in combination with erlotinib.

We have data with this combination in other tumors, and we know single-agent erlotinib has minor activity in recurrent ovarian cancer (Gordon 2005). Synergy may exist between these two biologic agents. In addition, high levels of EGFR expression occur in ovarian cancer cells.

We wouldn't expect erlotinib to do much for a patient with a large-volume recurrence. However, in the maintenance setting for a patient with small-volume disease, some of the anti-angiogenic effects of erlotinib might combine or perhaps synergize with bevacizumab to be beneficial.

Track 6

► **DR LOVE:** How are you using bevacizumab in your practice?

► **DR MATULONIS:** I don't use bevacizumab for patients with newly diagnosed ovarian cancer, but I do use it in recurrent disease. I select patients carefully and counsel them about the risks. I choose patients with a low risk of gastrointestinal perforation who are not hypertensive. I check their blood pressure and make sure they don't have any cardiovascular risk factors.

► **DR LOVE:** When the ascites in a patient with refractory, advanced disease responds to bevacizumab, is that an antitumor effect, a vascular effect or both?

► **DR MATULONIS:** When we see a reduction in ascites, we usually see concomitant decreases in peritoneal metastases, lymph node size and the areas of solid tumor growth. Although we can't be 100 percent certain, I believe it's an anticancer effect.

Track 9

► **DR LOVE:** What are your thoughts about anti-HER2 therapy in ovarian cancer, including the potential role of pertuzumab?

► **DR MATULONIS:** HER2 is not generally overexpressed in ovarian cancer, so we have to screen many cases to find the few that do overexpress HER2. Even in those patients with HER2 overexpression, the responses to trastuzumab are not particularly robust. Pertuzumab works differently than trastuzumab. Trastuzumab prevents HER2 signaling, while pertuzumab blocks HER2 receptor dimerization with HER3.

Three trials have evaluated pertuzumab in ovarian cancer. A single-agent study was published in the *Journal of Clinical Oncology* a number of years ago (Gordon 2006). A European Phase II trial evaluated a platinum-based chemotherapy regimen with or without pertuzumab in patients with platinum-sensitive disease (Kaye 2008). Another Phase II trial compared gemcitabine with or without pertuzumab in platinum-resistant ovarian cancer (Makhija 2007; [2.1]).

2.1

Phase II, Randomized, Placebo-Controlled, Double-Blind Trial of Gemcitabine with or without Pertuzumab for Patients with Platinum-Resistant Ovarian, Fallopian Tube or Primary Peritoneal Cancer

Efficacy parameters	Gemcitabine + placebo (n = 65)	Gemcitabine + pertuzumab (n = 65)
Progression-free survival		
Median (range)	2.6 months (0-12.7)	2.9 months (0-8.7)
Adjusted HR (95% CI)	0.66 (0.43-1.03)	
Rate at 4 months	37%	48%
Overall response rate	5%	14%
Survival		
Median (range)	13.1 months (1.5-25.4)	12.0 months (1.3-23.9)
Adjusted HR (95% CI)	0.99 (0.60-1.62)	
Time to symptom deterioration (FOSI)		
Median (range)	1.7 (0-10.2)	3.8 (0-7.7)
Adjusted HR (95% CI)	0.62 (0.36-1.05)	

HR = hazard ratio; CI = confidence interval; FOSI = Functional Assessment of Cancer Therapy-Ovarian (FACT-O) Symptom Index

SOURCE: Makhija S et al. *Proc ASCO* 2007; **Abstract 5507**.

 **Track 10**

▶ **DR LOVE:** Can you summarize the efficacy and toxicity associated with IP therapy?

▶ **DR MATULONIS:** In the GOG-172 trial, comparing IV paclitaxel/cisplatin to IV paclitaxel with IP cisplatin/paclitaxel in patients with Stage III disease, the improvement in overall survival of approximately 16 months in the IP arm was the best we have seen so far (Armstrong 2006; [2.2]). The GOG-111 study, which added IV paclitaxel to IV cisplatin and compared it to IV cisplatin/cyclophosphamide in patients with Stage III/IV ovarian cancer, reported an approximately one-year improvement in overall survival (McGuire 1996).

The downside of IP therapy is the toxicity, which is greater than that associated with IV regimens, including increased neuropathy, pancytopenia, nausea, vomiting, renal toxicities and electrolyte problems. The side effects are

ameliorated somewhat by reducing the cisplatin dose to 75 mg/m², but that dose hasn't been tested in a randomized manner, so it can't be recommended uniformly for use.

2.2

Intraperitoneal Cisplatin and Paclitaxel in Ovarian Cancer

“An intensive regimen of intravenous paclitaxel followed by intraperitoneal cisplatin and paclitaxel significantly improved progression-free survival ($P = 0.05$) and overall survival ($P = 0.03$) among women with newly diagnosed, optimally debulked stage III ovarian cancer. As compared with the intravenous-therapy group, women who received intraperitoneal treatment had a 25 percent reduction in the risk of death. Among all randomized phase 3 trials conducted by the GOG among patients with advanced ovarian cancer, the current trial yielded the longest median survival: 65.6 months, in the group of patients who received intraperitoneal therapy.”

SOURCE: Armstrong DK et al. *N Engl J Med* 2006;354(1):34-43. [Abstract](#)

Track 14

▶ **DR LOVE:** What is your clinical approach for patients with platinum-refractory ovarian cancer?

▶ **DR MATULONIS:** Outside of a clinical trial, treatment options include using one of several cytotoxic agents or bevacizumab. I usually choose either topotecan or pegylated liposomal doxorubicin (PLD). I do not use the FDA-approved dose of PLD but a lower dose — 40 mg/m². The lower dose is more tolerable for patients in terms of the rash. Generally, if the rash becomes a problem one can reduce the dose or stretch the interval between cycles. If the rash is Grade II or worse, I generally drop the dose from 40 to 32 mg/m². ■

SELECT PUBLICATIONS

Armstrong DK et al. **Intraperitoneal cisplatin and paclitaxel in ovarian cancer.** *N Engl J Med* 2006;354(1):34-43. [Abstract](#)

Gordon AN et al. **Efficacy and safety of erlotinib HCl, an epidermal growth factor receptor (HER1/EGFR) tyrosine kinase inhibitor, in patients with advanced ovarian carcinoma: Results from a phase II multicenter study.** *Int J Gynecol Cancer* 2005;15(5):785-92. [Abstract](#)

Gordon MS et al. **Clinical activity of pertuzumab (rhuMAb 2C4), a HER dimerization inhibitor, in advanced ovarian cancer: Potential predictive relationship with tumor HER2 activation status.** *J Clin Oncol* 2006;24(26):4324-32. [Abstract](#)

Kaye SB et al. **A randomised phase II study evaluating the combination of carboplatin-based chemotherapy with pertuzumab (P) versus carboplatin-based therapy alone in patients with relapsed, platinum sensitive ovarian cancer.** *Proc ASCO* 2008;[Abstract 5520](#).

Makhija S et al. **Results from a phase II randomized, placebo-controlled, double-blind trial suggest improved PFS with the addition of pertuzumab to gemcitabine in patients with platinum-resistant ovarian, fallopian tube, or primary peritoneal cancer.** *Proc ASCO* 2007;[Abstract 5507](#).

McGuire WP et al. **Cyclophosphamide and cisplatin compared with paclitaxel and cisplatin in patients with stage III and stage IV ovarian cancer.** *N Engl J Med* 1996;334(1):1-6. [Abstract](#)



INTERVIEW

Robert A Burger, MD

Dr Burger is Associate Professor of Clinical Obstetrics and Gynecology, Step II in the Department of Obstetrics and Gynecology's Division of Gynecologic Oncology at the University of California, Irvine Medical Center in Orange, California.

Tracks 1-12

- | | | | |
|----------------|--|-----------------|--|
| Track 1 | GOG-0170D: A Phase II study of bevacizumab in persistent or recurrent epithelial ovarian cancer or PPC | Track 8 | Continuation of bevacizumab after disease progression |
| Track 2 | Early clinical experience with bevacizumab in ovarian cancer | Track 9 | Caveats in the use of CA125 for monitoring disease progression in patients treated with anti-VEGF therapy |
| Track 3 | VEGF expression in ovarian cancer | Track 10 | Clinical algorithm for second- and later-line therapy in ovarian cancer |
| Track 4 | Activity of single-agent bevacizumab in recurrent ovarian cancer | Track 11 | Proposed Phase III trial of intraperitoneal chemotherapy in combination with bevacizumab for optimally debulked Stage III ovarian cancer |
| Track 5 | Gastrointestinal perforation events in Phase II trials of bevacizumab | Track 12 | Anticipated efficacy of adjuvant bevacizumab in ovarian cancer |
| Track 6 | Historical cohort studies of bevacizumab in heavily pretreated patients with ovarian cancer | | |
| Track 7 | Clinical use of single-agent bevacizumab in advanced ovarian cancer | | |

Select Excerpts from the Interview

Tracks 1, 4-5

► **DR LOVE:** Can you discuss the data from the GOG-0170D trial evaluating the use of bevacizumab for patients with persistent or recurrent epithelial ovarian cancer or primary peritoneal cancer?

► **DR BURGER:** Among 62 patients who had been previously treated with one or two cytotoxic regimens and had measurable disease by RECIST, bevacizumab led to a 21 percent objective response rate and a 40 percent progression-free survival of at least six months (Burger 2007; [3.1]).

A regression analysis performed at the conclusion of the trial factored in variables associated with disease progression in patients enrolled on Phase II trials of traditional cytotoxic agents: number of prior regimens, time from completion of initial therapy to first recurrence, age and performance status.

None of those factors predicted the time to disease progression in patients treated with bevacizumab (Burger 2007).

► **DR LOVE:** Can you compare GOG-0170D to the other recently reported Phase II trial of bevacizumab in patients with platinum-resistant ovarian cancer or peritoneal serous cancer (Cannistra 2007)?

► **DR BURGER:** In GOG-0170D, the patients had received either one or two prior regimens. About 40 of the patients had experienced disease progression within six months of receiving their most recent platinum-containing regimen, and the remainder had a platinum-free interval greater than or equal to six months (Burger 2007). The other single-agent bevacizumab trial required that the patients' disease be either primarily or secondarily platinum-resistant, and they could have received two to three prior regimens (Cannistra 2007).

So regarding risk of disease progression with any therapy, the patients in the trial reported by Cannistra were at higher risk. In the trial by Cannistra, the objective response rate was 16 percent (Cannistra 2007). In GOG-0170D, the response rate was 21 percent (Burger 2007; [3.1]).

► **DR LOVE:** What was observed in the two studies in terms of side effects and toxicity?

► **DR BURGER:** The differences in toxicity were interesting. It's hard to compare across trials, but the trial by Cannistra was closed prematurely due to five cases of gastrointestinal perforation, and GOG-0170D had zero cases of gastrointestinal perforation.

3.1

GOG-0170D: A Phase II Trial of Bevacizumab Monotherapy in Persistent or Recurrent Epithelial Ovarian Cancer or Primary Peritoneal Cancer (N = 62)

Efficacy data

Response rate	21% (90% CI: 12.9-31.3%)
Complete response	3.2%
Partial response	17.7%
Stable disease	51.6%
Median duration of response	10.3 months
Progression-free survival (PFS) \geq 6 months	40.3% (90% CI: 29.8-53.6%)

Conclusions

"In the second and third line treatment setting, patients with recurrent epithelial ovarian and primary peritoneal cancer, single agent bevacizumab:

- *Well tolerated at the dose and schedule of 15 mg/kg q21 days*
- *Active by clinical response and PFS"*

CI = confidence interval

SOURCE: Burger RA et al. *J Clin Oncol* 2007;25(33):5165-71. [Abstract](#)

In GOG-0170D, the rate of Grade III/IV proteinuria was minimal. Only one patient developed nephrotic syndrome, and it reversed after the discontinuation of bevacizumab. Approximately 10 percent of the patients had clinically relevant hypertension requiring antihypertensive therapy (Burger 2007). In the trial by Cannistra, no patients developed Grade III/IV proteinuria, but a number of patients with hypertension required therapy. Three patients experienced Grade III/IV arterial thrombotic events (Cannistra 2007).

Track 11

► **DR LOVE:** Can you discuss evolving clinical research on the use of bevacizumab in combination with IP therapy?

► **DR BURGER:** IP chemotherapy is considered a standard option for patients with Stage III ovarian cancer, especially those who have undergone optimal cytoreductive surgery. Three Phase III trials have demonstrated a survival advantage with IP regimens in combination with IV chemotherapy compared to standard IV chemotherapy regimens alone (Alberts 1996; Armstrong 2006; Markman 2001).

It's important to establish the safety and efficacy of these IP cytotoxic regimens in combination with bevacizumab. A Phase III trial is being developed to evaluate modified IP cytotoxic therapy in combination with bevacizumab for patients with Stage III, optimally debulked ovarian cancer. ■

SELECT PUBLICATIONS

Alberts DS et al. **Intraperitoneal cisplatin plus intravenous cyclophosphamide versus intravenous cisplatin plus intravenous cyclophosphamide for stage III ovarian cancer.** *N Engl J Med* 1996;335(26):1950-5. [Abstract](#)

Armstrong DK et al. **Intraperitoneal cisplatin and paclitaxel in ovarian cancer.** *N Engl J Med* 2006;354(1):34-43. [Abstract](#)

Burger RA et al. **Phase II trial of bevacizumab in persistent or recurrent epithelial ovarian cancer or primary peritoneal cancer: A Gynecologic Oncology Group study.** *J Clin Oncol* 2007;25(33):5165-71. [Abstract](#)

Cannistra SA et al. **Phase II study of bevacizumab in patients with platinum-resistant ovarian cancer or peritoneal serous cancer.** *J Clin Oncol* 2007;25(33):5180-6. [Abstract](#)

Duncan TJ et al. **Vascular endothelial growth factor expression in ovarian cancer: A model for targeted use of novel therapies?** *Clin Cancer Res* 2008;14(10):3030-5. [Abstract](#)

Hamilton CA et al. **Intraperitoneal bevacizumab for the palliation of malignant ascites in refractory ovarian cancer.** *Gynecol Oncol* 2008;[Epub ahead of print]. [Abstract](#)

Kaye SB. **Bevacizumab for the treatment of epithelial ovarian cancer: Will this be its finest hour?** *J Clin Oncol* 2007;25(33):5150-2. No abstract available

Markman M et al. **Phase III trial of standard-dose intravenous cisplatin plus paclitaxel versus moderately high-dose carboplatin followed by intravenous paclitaxel and intraperitoneal cisplatin in small-volume stage III ovarian carcinoma: An Intergroup study of the Gynecologic Oncology Group, Southwestern Oncology Group, and Eastern Cooperative Oncology Group.** *J Clin Oncol* 2001;19(4):1001-7. [Abstract](#)

Spannuth WA et al. **Angiogenesis as a strategic target for ovarian cancer therapy.** *Nat Clin Pract Oncol* 2008;5(4):194-204. [Abstract](#)



INTERVIEW

Bradley J Monk, MD

Dr Monk is Associate Professor and Director of Research in the Division of Gynecologic Oncology at the University of California, Irvine Medical Center's Chao Family Comprehensive Cancer Center in Orange, California.

Tracks 1-14

- Track 1** Primary treatment for patients with ovarian cancer
- Track 2** Standard therapy and controversies in the up-front treatment of epithelial ovarian cancer
- Track 3** Perspective on intraperitoneal cisplatin/paclitaxel
- Track 4** Current priority clinical trials in ovarian cancer: GOG-0218 (up-front chemotherapy/bevacizumab) and GOG-0212 (consolidation therapy)
- Track 5** Considerations for the clinical use of up-front chemotherapy with bevacizumab
- Track 6** Rationale for the effectiveness of anti-VEGF therapy in ovarian cancer
- Track 7** Clinical approach to patients with recurrent ovarian cancer
- Track 8** Phase III study of trabectedin with pegylated liposomal doxorubicin (PLD) versus PLD in relapsed recurrent ovarian cancer
- Track 9** Viewpoint on modest improvements in primary efficacy endpoints of clinical trials
- Track 10** Phase III study of patupilone versus PLD for taxane- or platinum-refractory/resistant recurrent epithelial ovarian cancer, PFC or PPC
- Track 11** Randomized Phase II study of gemcitabine with or without pertuzumab in ovarian cancer
- Track 12** HER pathway gene expression analysis and prediction of benefit from pertuzumab
- Track 13** Timing of initiation and discontinuation of treatment for recurrent ovarian cancer
- Track 14** Bevacizumab-associated side effects during treatment of recurrent ovarian cancer

Select Excerpts from the Interview

Track 2

► **DR LOVE:** What are some of the questions medical oncologists and gynecologic oncologists ask about the management of newly diagnosed ovarian cancer?

► **DR MONK:** The standard up-front treatment for epithelial ovarian cancer is maximal surgical debulking followed by six courses of intravenous platinum- and taxane-based chemotherapy administered every three weeks. I believe that

guidelines have established the platinum drug as carboplatin at an area under the curve (AUC) between 5.0 and 7.5 mg/mL and the taxane as paclitaxel at a dose of 175 mg/m² administered over three hours (NCCN 2008).

However, questions then emanate and alter that standard. One can outline seven acceptable modifications: (1) using IP chemotherapy, (2) administering more than six cycles of chemotherapy, (3) using weekly chemotherapy, (4) substituting docetaxel as the taxane, (5) adding a targeted agent, specifically bevacizumab, (6) using reassessment surgery when the disease is in remission at the completion of adjuvant chemotherapy to confirm whether the tumor is in remission and (7) continuing chemotherapy during remission as maintenance or consolidation therapy.

Track 4

► **DR LOVE:** What clinical trials are ongoing for patients in the up-front treatment setting?

► **DR MONK:** One of the two scientific priorities being evaluated in the up-front treatment of epithelial ovarian cancer is the incorporation of bevacizumab. The GOG-0218 trial (4.1) adds bevacizumab to a platinum-and-taxane backbone. It also addresses maintenance bevacizumab in a third arm, even when the patient's disease is in remission.


4.1

GOG-0218: A Phase III Study of Carboplatin/Paclitaxel (CP) versus CP/Bevacizumab with or without Extended Bevacizumab

Protocol IDs: GOG-0218, NCT00262847; Target Accrual: 2,000 (Open)

Eligibility

- Histologically confirmed Stage III with any gross residual disease OR Stage IV epithelial ovarian, primary peritoneal or fallopian tube cancer
- No prior chemotherapy for abdominal or pelvic cancer
- At least four weeks since major surgical procedure or open biopsy



¹ Chemotherapy = (IV paclitaxel 175 mg/m² + IV carboplatin AUC 6 mg/mL x min) every 21 days x 6 cycles

² Bevacizumab = 15 mg/kg every 21 days (beginning cycle 2)

SOURCES: NCI Physician Data Query, November 2008; Burger RA et al. *Proc ASCO* 2005; **Abstract 5009**.

The second scientific priority, also one of my seven controversies, is consolidation therapy. GOG-0212 (4.2) randomly assigns patients who are in remission to no treatment, 12 months of paclitaxel or 12 months of polyglutamate paclitaxel.

4.2

GOG-0212: A Phase III Randomized Trial of Paclitaxel versus Polyglutamate Paclitaxel as Consolidation/Maintenance Therapy versus No Further Anticancer Therapy until Disease Progression

Target Accrual: 1,100 (Open)

Eligibility

- Stage III/IV ovarian epithelial or primary peritoneal cancer
- Surgery
- Primary therapy: Five to eight courses of carboplatin (IV or IP) and paclitaxel or docetaxel



IV = intravenous; IP = intraperitoneal

SOURCE: NCI Physician Data Query, November 2008.

Track 8

▶ **DR LOVE:** Could you discuss your research on trabectedin?

▶ **DR MONK:** At the 2008 European Society of Medical Oncology meeting, I presented a trial with trabectedin — a DNA-active drug initially derived from the Caribbean sea squirt. I evaluated PLD with or without trabectedin in almost 700 patients. The combination demonstrated an improvement in

4.3

Randomized Phase III Trial of Trabectedin with Pegylated Liposomal Doxorubicin (PLD) versus PLD Alone in Women (N = 672) with Relapsed Recurrent Ovarian Cancer

	PLD + trabectedin	PLD alone	Hazard ratio	p-value
Median PFS				
All patients	7.3 months	5.8 months	0.79	0.019
Patients with PFI > 6 months	9.2 months	7.5 months	0.73	0.017
Response rate				
All patients	28%	19%	—	0.008
Patients with PFI > 6 months	35%	23%	—	0.004

PFS = progression-free survival; PFI = platinum-free interval

SOURCE: Monk BJ et al. ESMO Congress 2008; [Abstract LBA4](#).

progression-free survival and response rate. Most of the benefit was observed in patients with chemotherapy-sensitive disease (Monk 2008; [4.3]).

Phase II trials of single-agent trabectedin have shown substantial activity in patients with chemotherapy-sensitive disease, with response rates between 30 and 40 percent. In patients with chemotherapy-resistant disease, the response rates were between five and 10 percent (Krasner 2007; Sessa 2005).

Tracks 11-12

▶ **DR LOVE:** Would you discuss what you know about pertuzumab?

▶ **DR MONK:** Pertuzumab is interesting because it is an antibody to HER3, which seems to be an important biomarker and target in epithelial ovarian cancer. Pertuzumab activity was demonstrated in the report of a randomized Phase II study of gemcitabine with or without pertuzumab (Makhija 2007; [2.1, page 9]). The final data from that study reported at ASCO 2008 suggested an association between efficacy and HER3 gene expression levels by RT-PCR (Amler 2008).

That endpoint was exploratory, so we question whether the evidence is sufficient or if we need to validate HER3 as a biomarker in a large prospective study before proceeding with studies of pertuzumab based on that biomarker. We need to study these agents in settings in which they will have the greatest likelihood of being effective, so I believe we should attempt to enrich the patient populations. We should use HER3 as a biomarker now to study pertuzumab. If the results are positive, the study should expand to a broader population. ■

SELECT PUBLICATIONS

Amler L et al. **HER pathway gene expression analysis in a phase II study of pertuzumab + gemcitabine vs gemcitabine + placebo in patients with platinum-resistant epithelial ovarian cancer.** ASCO 2008; [Abstract 5552](#).

Burger RA et al. **Phase II trial of bevacizumab in persistent or recurrent epithelial ovarian cancer (EOC) or primary peritoneal cancer (PPC): A Gynecologic Oncology Group (GOG) study.** *Proc ASCO* 2005; [Abstract 5009](#).

Krasner CN et al. **A Phase II study of trabectedin single agent in patients with recurrent ovarian cancer previously treated with platinum-based regimens.** *Br J Cancer* 2007;97(12):1618-24. [Abstract](#)

Makhija S et al. **Results from a phase II randomized, placebo-controlled, double-blind trial suggest improved PFS with the addition of pertuzumab to gemcitabine in patients with platinum-resistant ovarian, fallopian tube, or primary peritoneal cancer.** ASCO 2007; [Abstract 5507](#).

Monk BJ et al. **A randomized phase III study of trabectedin with pegylated liposomal doxorubicin (PLD) versus PLD in relapsed, recurrent ovarian cancer.** ESMO Congress 2008; [Abstract LBA4](#).

National Comprehensive Cancer Network (NCCN®). **NCCN clinical practice guidelines in oncology. Ovarian cancer — Version 1. 2008.** Available at: http://nccn.org/professionals/physician_gls/PDF/ovarian.pdf.

Sessa C et al. **Trabectedin for women with ovarian carcinoma after treatment with platinum and taxanes fails.** *J Clin Oncol* 2005;23(9):1867-74. [Abstract](#)

QUESTIONS (PLEASE CIRCLE ANSWER):

1. In an ongoing Phase II trial at the Dana-Farber/Harvard Cancer Center, patients with ovarian cancer receive IV carboplatin/paclitaxel with bevacizumab for six months and then are randomly assigned to receive bevacizumab with or without _____.
 - a. Cetuximab
 - b. Erlotinib
 - c. Pertuzumab
2. Which agent is believed to inhibit HER dimerization?
 - a. Cetuximab
 - b. Pertuzumab
 - c. Trastuzumab
3. In GOG-172, comparing IV paclitaxel/cisplatin to IV paclitaxel with IP cisplatin/paclitaxel for patients with Stage III disease, overall survival was approximately _____ months longer for the patients receiving IP therapy.
 - a. Two
 - b. Six
 - c. 12
 - d. 16
4. In a Phase II trial (GOG-0170D) for patients with persistent or recurrent epithelial ovarian cancer or primary peritoneal cancer, bevacizumab monotherapy resulted in a response rate of approximately _____.
 - a. Two percent
 - b. Seven percent
 - c. 21 percent
 - d. 40 percent
5. In GOG-0170D, which evaluated bevacizumab monotherapy for patients with recurrent ovarian cancer or primary peritoneal cancer, no cases of bowel perforations were reported.
 - a. True
 - b. False
6. GOG-0218 will evaluate the role of _____ in combination with paclitaxel/carboplatin for women with newly diagnosed Stage III/IV epithelial ovarian, primary peritoneal or fallopian tube cancer.
 - a. Bevacizumab
 - b. Pertuzumab
 - c. Trastuzumab
 - d. Gemcitabine
 - e. None of the above
7. GOG-0212 will evaluate the role of _____ as consolidation/maintenance therapy for women with Stage III/IV epithelial ovarian or primary peritoneal cancer who have been treated with surgery and five to eight courses of primary chemotherapy.
 - a. Paclitaxel
 - b. Polyglutamate paclitaxel
 - c. Bevacizumab
 - d. Both a and b
 - e. Both a and c
8. Which of the following agents has received FDA approval for the treatment of recurrent ovarian cancer?
 - a. Topotecan
 - b. PLD
 - c. Bevacizumab
 - d. Both a and b
 - e. All of the above
9. The addition of trabectedin to _____ was found to improve progression-free survival and response rates for women with recurrent ovarian cancer.
 - a. Bevacizumab
 - b. Topotecan
 - c. PLD
 - d. All of the above
 - e. None of the above

Research To Practice is committed to providing valuable continuing education for oncology clinicians, and your input is critical to helping us achieve this important goal. Please take the time to assess the activity you just completed, with the assurance that your answers and suggestions are strictly confidential.

PART ONE — Please tell us about your experience with this educational activity

BEFORE completion of this activity, how would you characterize your level of knowledge on the following topics?

4 = Very good 3 = Above average 2 = Adequate 1 = Suboptimal

Management of platinum-resistant, recurrent ovarian cancer	4	3	2	1
Toxicities associated with intraperitoneal therapy	4	3	2	1
Role of anti-angiogenic and other targeted agents in the treatment of ovarian cancer	4	3	2	1
Rationale for maintenance therapy in optimally debulked, Stage II/III disease	4	3	2	1

AFTER completion of this activity, how would you characterize your level of knowledge on the following topics?

4 = Very good 3 = Above average 2 = Adequate 1 = Suboptimal

Management of platinum-resistant, recurrent ovarian cancer	4	3	2	1
Toxicities associated with intraperitoneal therapy	4	3	2	1
Role of anti-angiogenic and other targeted agents in the treatment of ovarian cancer	4	3	2	1
Rationale for maintenance therapy in optimally debulked, Stage II/III disease	4	3	2	1

Was the activity evidence based, fair, balanced and free from commercial bias?

Yes No

Please explain:

Will this activity help you improve patient care?

Yes No Not applicable

If no, please explain:

Did the activity meet your educational needs and expectations?

Yes No

If no, please explain:

Please respond to the following LEARNER statements by circling the appropriate selection:

4 = Yes 3 = Will consider 2 = No 1 = Already doing N/M = Learning objective not met N/A = Not applicable

As a result of this activity, I will be able to:

- Recognize the contributory roles of the specialized interdisciplinary team in achieving best-practice surgical and medical outcomes for patients with ovarian cancer4 3 2 1 N/M N/A
- Compare and contrast the risks and benefits of intraperitoneal and intravenous chemotherapy regimens when devising management strategies for patients with optimally debulked Stage II and Stage III ovarian cancer4 3 2 1 N/M N/A
- Develop an evidence-based algorithm for the systemic treatment of recurrent platinum-sensitive and platinum-resistant ovarian cancer that optimizes long-term patient outcome and quality of life.....4 3 2 1 N/M N/A
- Summarize the existing data and ongoing clinical trials focused on angiogenesis inhibition of ovarian cancer, and identify patients who may benefit from this therapeutic approach4 3 2 1 N/M N/A
- Consider the utility of evaluating CA125 serum levels and radiographic monitoring of patients with ovarian cancer that is in a state of remission4 3 2 1 N/M N/A
- Understand the potential role of the risk-adapted use of maintenance systemic therapy for patients with ovarian cancer who have demonstrated an initial treatment response4 3 2 1 N/M N/A
- Recall the scientific rationale for molecular-targeted agents under active investigation for the treatment of ovarian cancer4 3 2 1 N/M N/A
- Counsel appropriately selected patients with ovarian cancer about the availability of and participation in ongoing clinical trials.....4 3 2 1 N/M N/A

EDUCATIONAL ASSESSMENT AND CREDIT FORM (continued)

What other practice changes will you make or consider making as a result of this activity?

What additional information or training do you need on the activity topics or other oncology-related topics?

Additional comments about this activity:

As part of our ongoing, continuous quality-improvement effort, we conduct postactivity follow-up surveys to assess the impact of our educational interventions on professional practice. Please indicate your willingness to participate in such a survey.

Yes, I am willing to participate in a follow-up survey. No, I am not willing to participate in a follow-up survey.

PART TWO — Please tell us about the editor and faculty for this educational activity

	4 = Very good	3 = Above average	2 = Adequate	1 = Suboptimal	
Faculty					
	Knowledge of subject matter				Effectiveness as an educator
Tate Thigpen, MD	4	3	2	1	4 3 2 1
Ursula A Matulonis, MD	4	3	2	1	4 3 2 1
Robert A Burger, MD	4	3	2	1	4 3 2 1
Bradley J Monk, MD	4	3	2	1	4 3 2 1
Editor					
	Knowledge of subject matter				Effectiveness as an educator
Neil Love, MD	4	3	2	1	4 3 2 1

Please recommend additional faculty for future activities:

Other comments about the editor and faculty for this activity:

REQUEST FOR CREDIT — Please print clearly

Name: Specialty:

Professional Designation:

MD DO PharmD NP RN PA Other

Medical License/ME Number: Last 4 Digits of SSN (required):

Street Address: Box/Suite:

City, State, Zip:

Telephone: Fax:

Email:

Research To Practice designates this educational activity for a maximum of 3 AMA PRA Category 1 Credits™. Physicians should only claim credit commensurate with the extent of their participation in the activity.

I certify my actual time spent to complete this educational activity to be _____ hour(s).

Signature: Date:

To obtain a certificate of completion and receive credit for this activity, please complete the Post-test, fill out the Educational Assessment and Credit Form and fax both to (800) 447-4310, or mail both to Research To Practice, One Biscayne Tower, 2 South Biscayne Boulevard, Suite 3600, Miami, FL 33131. You may also complete the Post-test and Educational Assessment online at www.ResearchToPractice.com/OCU/CME.

Ovarian Cancer™

U P D A T E

Editor	Neil Love, MD
Managing Editor	Kathryn Ault Ziel, PhD
Scientific Director	Richard Kaderman, PhD
Senior Director, Medical Affairs	Aviva Asnis-Alibozek, PA-C, MPAS
Writers	Lillian Sklaver Poltorack, PharmD Douglas Paley Clayton Campbell Anne Jacobson, MPH
Continuing Education Administrator for Nursing	Sally Bogert, RNC, WHCNP
Content Validation	Margaret Peng Erin Wall Jessica McCarrick
Director, Creative and Copy Editing	Aura Herrmann
Creative Manager	Fernando Rendina
Graphic Designers	Jessica Benitez Jason Cunnius Tamara Dabney Shantia Daniel Claudia Munoz
Senior Production Editor	Alexis Oneca
Traffic Manager	Tere Sosa
Copy Editors	Dave Amber Margo Harris David Hill Rosemary Hulce Kirsten Miller Pat Morrissey/Havlin Carol Peschke Susan Petrone
Production Manager	Tracy Potter
Audio Production	Frank Cesarano
Web Master	John Ribeiro
Faculty Relations Manager	Melissa Vives
CME Director/CPD Director	Isabelle Tate
Contact Information	Neil Love, MD Research To Practice One Biscayne Tower 2 South Biscayne Boulevard, Suite 3600 Miami, FL 33131 Fax: (305) 377-9998 Email: DrNeilLove@ResearchToPractice.com Email: CE@ResearchToPractice.com
For CME/CNE Information	

Copyright © 2008 Research To Practice. All rights reserved.

The compact discs, Internet content and accompanying printed material are protected by copyright. No part of this program may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or utilizing any information storage and retrieval system, without written permission from the copyright owner.

The opinions expressed are those of the presenters and are not to be construed as those of the publisher or grantors.

Participants have an implied responsibility to use the newly acquired information to enhance patient outcomes and their

own professional development. The information presented in this activity is not meant to serve as a guideline for patient management.

Any procedures, medications or other courses of diagnosis or treatment discussed or suggested in this activity should not be used by clinicians without evaluation of their patients' conditions and possible contraindications or dangers in use, review of any applicable manufacturer's product information and comparison with recommendations of other authorities.

Ovarian Cancer™

U P D A T E

Copyright © 2008 Research To Practice.
This program is supported by educational grants from Eli Lilly and Company,
Genentech BioOncology and Ortho Biotech Products LP.

Research
To Practice®

Sponsored by Research To Practice.

Last review date: December 2008
Release date: December 2008
Expiration date: December 2009
Estimated time to complete: 3 hours